

Ketamine이 청각 기억에 미치는 영향 : Mismatch Negativity와 정신분열병의 Glutamate 가설

윤 탁 · 안경흡 · 배지수 · 김명선 · 김재진 · 권준수

ABSTRACT

Effect of Ketamine on the Echoic Memory Process : The Mismatch Negativity and Glutamate Receptor System in Schizophrenia

Tak Youn, MD, Kyung Heup Ahn, MD, Ji Soo Pae, MD,
Myung Sun Kim, PhD, Jae-Jin Kim, MD, PhD and Jun Soo Kwon, MD, PhD
Department of Psychiatry, Seoul National University College of Medicine, Seoul, Korea

Objective : The abnormality of mismatch negativity (MMN) in schizophrenia is thought to be associated with perceptual disturbance and cognitive dysfunction. And the antagonists of the N-methyl-D-aspartate (NMDA) receptors, ketamine, can induce anomalies of psychophysiology and cognitive function as those of schizophrenia. In order to explore the role of NMDA receptors on echoic memory system, MMN under ketamine administration was analyzed. **Methods** : MMNs of Healthy 12 subjects under sub-anesthetic dose (0.65 mg/kg/hr) of ketamine administration in placebo-controlled design were recorded by 128 channel EEG. Brief Psychiatric Rating Scale (BPRS) change was also evaluated. **Results** : BPRS score was significantly increased by ketamine administration ($t = -6.655, p < 0.001$). Ketamine induced significant decrease in MMN amplitudes ($F_z, t = -2.572, p = 0.026$). Neither MMN amplitude under placebo administration nor MMN latencies under ketamine administration and placebo was changed significantly. **Conclusion** : Ketamine induced echoic memory dysfunction in healthy subjects, which is usually found in schizophrenic patients. Consequently, reduced glutamatergic activity in brain could be involved some early processes of the memory dysfunction in schizophrenia. **(Korean J Psychopharmacol 2001;12(4):322-327)**

KEY WORDS : Schizophrenia Information process · NMDA receptor · Mismatch negativity · Echoic memory.

서 론

(04 - 1999 - 055 - 0)

: , 110 - 744

28

: (02) 760 - 2972 ·

: (02) 747 - 9063

E - mail : kwonjs@plaza.snu.ac.kr

Ketamine

(single-blind method)
ketamine(0.65 mg/kg/hr)
()
(1). ketamine
(Brief Psychiatric Rating Scale, BPRS)²³⁾
(baseline)
0.65 mg/kg/hr ketamine
(subanesthetic dosage)
(24-26)
(100 cc/hr)
BPRS MMN
ketamine
ketamine
0.26
mg/kg ketamine (bolus in -
jection) , ketamine 0.65 rate)
mg/kg/hr
ketamine 가
10
BPRS , MMN epoch
가
, ketamine (age)
) 15 50 msec MMN . MMN
, 2 30 BPRS
가 100 200 msec
MMN . MMN




Fig. 1. Summary of the procedure to get EEG under subanesthetic dosage of ketamine or placebo.

2) 청각 Oddball paradigm의 구성
STIM (Neurosoft Inc. USA)
1600
1520 (95%,
1000 Hz, 80 dB) 80 (5%, 1200 Hz,
80 dB) , oddball paradigm

100 msec, - 10 msec,

Table 1. The changes of the Brief Psychotic Rating Scale(BPRS) scores after ketamine and placebo injections

Medication	BPRS Scores (Mean \pm S.D.)
Ketamine* Pre-injection	1.00 \pm 0.74
Post-injection	6.58 \pm 2.64
Placebo Pre-injection	1.16 \pm 0.83
Post-injection	0.75 \pm 0.75

* : paired t-test, $p < 0.001$




Fig. 2. Effect of ketamine and placebo injection on mismatch negativity(MMN). MMNs were shown mainly in frontal area. Amplitude of MMN under ketamine administration was decreased significantly ($p < 0.05$).

Table 2. The changes of amplitude and latency of mismatch negativity after injections of ketamine or placebo

Medication	Amplitude(Fz, V)	Latency(Fz, msec)
Ketamine Pre-injection	- 5.73 \pm 2.34*	147.9 \pm 26.8
Post-injection	- 4.22 \pm 1.59*	149.3 \pm 22.1
Placebo Pre-injection	- 3.66 \pm 2.36	139.0 \pm 20.9
Post-injection	- 3.91 \pm 2.46	141.0 \pm 24.6

* : paired t-test, $p = 0.026$

300 msec

3) 결과 분석

MMN 가 Fz MMN
Ketamine
BPRS paired t - test
($p < 0.05$) SPSS
10.0(SPSS Inc, USA)

결 과

12 28 (SD = 2.30)
17.7
Ketamine BPRS
($t = - 6.655$, $p < 0.001$),
가 ($t = 1.449$, $p = 0.175$)(1).
MMN 2 Ketamine
, Fz 가
(Fz, $t = - 2.572$, $p = 0.026$).
가 keta -
mine 가
(2).

고 찰

MMN

12,21,28 - 31)

MMN

28) 12)

32)

Javitt MMN glutamate ketamine ket -

NMDA 20-22) amine .

MN NMDA Umricht MN 가 , Oranje MN 가

(0.9 mg/kg/hr)²⁰⁾ (0.9 mg/g kg/hr)²¹⁾

ketamine(0.9 mg/kg/hr) 0.65 mg/kg/hr ketamine BPRS ketamine

MMN ketamine , MMN ketamine

NMDA ketamine , NMDA MMN 가 , NMDA

keta mine PCP/NMDA 가

g ketamine echoic memory process)

가 , MMN (12) ,)

0.3 0.65 mg/kg/hr , NMDA 28)

0.65 mg/kg/hr ketamine 중심 단어 : Ketamine Mismatch nega-

Krystal 34) tivity

BPRS 참 고 문 헌

BPRS ke -

tamine 가

Malhotra 27,35) Krystal 34) 6 9

BPRS

ke -

참고문헌

1) Goldman-Rakic PS. Working memory dysfunction in schizophrenia. J Neuropsychiatry Clin Neurosci 1994;6:348-357.

2) Grotta J, Clark W, Coull B, Pettigrew LC, Mackay B, Goldstein LB, et al. Safety and tolerability of the glutamate antagonist CGS 1975(Selfotel) in patients with acute ischemic stroke. Results of a phase IIa randomized trial. Stroke 1995;26:602-605.

3) Kim JS, Kornhuber HH, Holzmüller B, Schmid-Burgk W, Mergner T, Krzepinski G. Reduction of cerebrospinal fluid

- glutamic acid in Huntington's chorea and in schizophrenic patients. *Arch Psychiatr Nervenkr* 1980;228:7-10.
- 4) Lipska BK, Weinberger DR. To model a psychiatric disorder in animals: Schizophrenia as a reality test. *Neuropsychopharmacology* 2000;23:223-239.
 - 5) Ross RG, Olincy A, Harris JG, Sullivan B, Radant A. Smooth pursuit eye movements in schizophrenia and attentional dysfunction: Adults with schizophrenia, ADHD, and a normal comparison group. *Biol Psychiatry* 2000;48:197-203.
 - 6) Tsai G, Passani LA, Slusher BS, Carter R, Baer L, Kleinman JE, et al. Abnormal excitatory neurotransmitter metabolism in schizophrenic brains. *Arch Gen Psychiatry* 1995;52:829-836.
 - 7) Cotman CW, Monaghan DT, Ganong AH. Excitatory amino acid neurotransmission: NMDA receptors and Hebb-type synaptic plasticity. *Annu Rev Neurosci* 1998;11:61-80.
 - 8) Näätänen R. The mismatch negativity. In: Näätänen R, editor. *Attention and Brain Function*. Hillsdale, NJ: Lawrence Erlbaum Associates;1992. p.136-200.
 - 9) Javitt DC. Intracortical mechanisms of mismatch negativity dysfunction in schizophrenia. *Audiol Neurotol* 2000;5:207-215.
 - 10) Shelley AM, Ward PB, Catts SV, Michie PT, Andrews S, McConaghy N. Mismatch negativity: An index of a preattentive processing deficit in schizophrenia. *Biol Psychiatry* 1991;30:1059-1062.
 - 11) Hirayasu Y, Potts GF, O'Donnell BF, Kwon JS, Arakaki H, Akdag SJ, et al. Auditory mismatch negativity in schizophrenia: Topographic evaluation with a high-density recording montage. *Am J Psychiatry* 1998;155:1281-1284.
 - 12) Youn T, Park HJ, Kim JJ, Kim MS, Kwon JS. Altered Hemispheric Asymmetry and Positive Symptoms in Schizophrenia: Equivalent Current Dipole of Auditory Mismatch Negativity. *Schizophr Res (in press)*.
 - 13) Scherg M, Vajsar J, Picton TW. A source analysis of the late human auditory evoked potentials. *J Cogn Neurosci* 1989;1:336-355.
 - 14) Wible CG, Kubicki M, Yoo SS, Kacher DF, Salisbury DF, Anderson MC, et al. A functional magnetic resonance imaging study of auditory mismatch in schizophrenia. *Am J Psychiatry* 2001;158:938-943.
 - 15) Barta PE, Pearlson GD, Powers RE, Richards SS, Tune LE. Auditory hallucinations and smaller superior temporal gyrus volume in schizophrenia. *Am J Psychiatry* 1990;147:1457-1462.
 - 16) Gur RE, Turetsky BI, Cowell PE, Finkelman C, Maany V, Grossman RI, et al. Temporolimbic volume reductions in schizophrenia. *Arch Gen Psychiatry* 2000;57:769-775.
 - 17) Kwon JS, McCarley RW, Hirayasu Y, Anderson JE, Fischer IA, Kikinis R, et al. Left planum temporale volume reduction in schizophrenia. *Arch Gen Psychiatry* 1999;56:142-148.
 - 18) Shenton ME, Kikinis R, Jolesz FA, Pollak SD, LeMay M, Wible CG, et al. Abnormalities of the left temporal lobe and thought disorder in schizophrenia. A quantitative magnetic resonance imaging study. *N Engl J Med* 1992;327:604-612.
 - 19) Kreitschmann-Andermahr I, Rosburg T, Meier T, Volz HP, Nowak H, Sauer H. Impaired sensory processing in male patients with schizophrenia: A magnetoencephalographic study of auditory mismatch detection. *Schizophr Res* 1999;35:121-129.
 - 20) Oranje B, van Berckel BN, Kemner C, van Ree JM, Kahn RS, Verbaten MN. The effects of a sub-anaesthetic dose of ketamine on human selective attention. *Neuropsychopharmacology* 2000;22:293-302.
 - 21) Umbricht D, Schmid L, Koller R, Vollenweider FX, Hell D, Javitt DC. Ketamine-induced deficits in auditory and visual context-dependent processing in healthy volunteers: Implications for models of cognitive deficits in schizophrenia. *Arch Gen Psychiatry* 2000;57:1139-1147.
 - 22) Javitt DC, Steinschneider M, Schroeder CE, Arezzo JC. Role of cortical N-methyl-D-aspartate receptors in auditory sensory memory and mismatch negativity generation: Implications for schizophrenia. *Proc Natl Acad Sci U S A* 1996;93:11962-11967.
 - 23) Overall JE, Gorham DR. The brief psychiatric rating scale. *Psychol Rep* 1962;10:799-812.
 - 24) Hedlund JL, Vieweg BW. The brief psychiatric rating scale (BPRS): A comprehensive review. *J Oper Psychiatr* 1980;11:48-64.
 - 25) Corssen G, Dommino EF. Dissociative anesthesia: Further pharmacologic studies and first clinical experiences with phencyclidine derivative CI-581. *Anesth Analg* 1966;45:29-40.
 - 26) Krystal JH, Karper LP, Seibyl JP, Freeman GK, Delaney R, Bremner JD, et al. Subanesthetic effects of the noncompetitive NMDA antagonist, ketamine, in humans. *Arch Gen Psychiatry* 1994;51:199-214.
 - 27) Malhotra AK, Pinals DA, Adler CM, Elman I, Clifton A, Pic-kar D, et al. Ketamine-induced exacerbation of psychotic symptoms and cognitive impairment in neuroleptic-free schizophrenics. *Neuropsychopharmacology* 1997a;17:141-150.
 - 28) Catts SV, Shelley AM, Ward PB, Liebert B, McConaghy N, Andrews S, et al. Brain potential evidence for an auditory sensory memory deficit in schizophrenia. *Am J Psychiatry* 1995;152:213-219.
 - 29) Javitt DC, Doneshka P, Grochowski S, Ritter W. Impaired mismatch negativity generation reflects widespread dysfunction of working memory in schizophrenia. *Arch Gen Psychiatry* 1995;52:550-558.
 - 30) Kasai K, Okazawa K, Nakagome K, Hiramatsu KI, Hata A, Fukuda M, et al. Mismatch negativity and N2b attenuation as an indicator for dysfunction of the preattentive and controlled processing for deviance detection in schizophrenia: A topographic event-related potential study. *Schizophr Res* 1999;35:141-156.
 - 31) Shelley AM, Silipo G, Javitt D. Diminished responsiveness of ERPs in schizophrenia subjects to changes in auditory stimulation parameters: Implications for theories of cortical dysfunction. *Schizophr Res* 1999;37:65-70.
 - 32) Oades RD, Dittmann-Balcar A, Zerbin D, Grzella I. Impaired attention dependent augmentation of MMN in nonparanoid vs. paranoid schizophrenic patients: A comparison with obsessive-compulsive disorder and healthy subjects. *Biol Psychiatry* 1997;41:1196-1210.
 - 33) Anand A, Charney DS, Oren DA, Berman RM, Hu XS, Capiello A, et al. Attenuation of the neuropsychiatric effects of ketamine with lamotrigine: Support for hyperglutamatergic effects of N-methyl-D-aspartate receptor antagonists. *Arch Gen Psychiatry* 2000;57:270-276.
 - 34) Krystal JH, Karper LP, Bennett A, D'Souza DC, Abi-Dargham A, Morrissey K, et al. Interactive effects of subanesthetic ketamine and subhypnotic lorazepam in humans. *Psychopharmacology* 1998;135:213-229.
 - 35) Malhotra AK, Adler CM, Kennison SD, Elman I, Pickar D, Breier A. Clozapine blunts N-methyl-D-aspartate antagonist-induced psychosis: A study with ketamine. *Biol Psychiatry* 1997b;42:664-668.
 - 36) Newcomer JW, Farber NB, Jevtovic-Todorovic V, Selke G, Melson AK, Hershey T, et al. Ketamine-induced NMDA receptor hypofunction as a model of memory impairment and psychosis. *Neuropsychopharmacology* 1999;20:106-118.